### **Ecoleaf Environmental Labeling Program**

Sustainable Management Promotion Organization 2-1, Kaji-cho 2 chome, Chiyoda-ku, Tokyo Japan https://ecoleaf-label.jp/

Type III Environmental Declaration (EPD)

Registration number: JR-BC-20002E

## Nakamoto Zourin Co.,Ltd

## Yakisugi [Shou Sugi Ban] / Gendai



ブラシ (G e n d a i )



Gendai施工例(米国)



Gendai施工例(日本)

### **Functional unit**

1 m<sup>2</sup> (15mm thick)

### **System boundary**

- ☐ final products
- intermediate products
- Inclusive of: A1 Raw Material Supply, A2 Transport, A3 Manufacturing
- Exclusive of: A4 Transport, A5 Construction, B1 Use, B2 Maintenance, B3 Repair, B4 Replacement, B5 Refurbishment, B6 Operational energy use, B7 Operational water use, C1 Demolition, C2 Transport, C3 Waste processing, C4 Disposal

### Main specifications of the product

Weight: 6.1kg/m2No paint applied

- Production sites: Hiroshima and Tokushima

### **Company Information**

Nakamotozourin Co.,Ltd https://nakamotozourin.co.jp Nakamoto Forestry North America https://nakamotoforestry.com Nakamoto Forestry Europe https://nakamotoforestry.eu

F	Registration#	JR-BC-20002E	
	PCR number	PA-120000-BC-01	
	PCR name	Wood、WoodMaterials	
P	ublication date	04/06/2020	
Ve	erification date	03/12/2020	
Ve	rification method	Product-by-product	
	Verification#	JR-BC-20002	
E	xpiration date	03/12/2025	
PC	PCR review was conducted by:		
	Approval date	12/25/2019	
	PCR review	Masayuki Kanzaki	
	panel chair	(Sustainable Management Promotion Organization)	

Tomoko Fuchigami

Independent verification of data & declaration in accordance with ISO14025

□internal

■ external

Registration number: JR-BC-20002E

<sup>\*</sup>Auditor's name is stated if system certification has been performed.

# EcoLeaf EcoLeaf

Acidification

**Ecoleaf Environmental Labeling Program** 


Sustainable Management Promotion Organization 2-1, Kaji-cho 2 chome, Chiyoda-ku, Tokyo Japan https://ecoleaf-label.jp/

# Type III Environmental Declaration (EPD) Registration number: JR-BC-20002E

1. Results of life cycle impact assessment (LCIA 0%

Global warming IPCC2013 GWP100a 2.2 kg-CO2eq

Ozone layer destruction 0.000018 g-CFC-11eq



Stage Parameter	Unit	Total	A1 Raw Material Supply	A2 Transport	A3 Manufacturin g	
Global warming IPCC2013 GWP100a	kg-CO <sub>2</sub> eq	2.2E+00	3.6E-01	3.6E-01	1.4E+00	
Ozone layer destruction	kg-CFC-11eq	1.8E-08	1.8E-08	2.9E-12	8.5E-12	
Acidification	kg-SO <sub>2</sub> eq	1.7E-03	1.6E-04	9.2E-04	6.5E-04	
Photochemical ozone	kg-C <sub>2</sub> H₄eq	2.9E-05	1.3E-06	9.3E-06	1.9E-05	
Eutrophication	kg-PO <sub>4</sub> 3-eq	5.8E-06	5.8E-06	2.5E-15	9.4E-15	


1.7 g-SO2eq

### 2. Life cycle inventory analysis (LCI) Unit **Parameter** Non-renewable material resources kg 1.3E-02 Non-renewable energy resources kg 7.5E-01 Non-renewable energy resources MJ 3.3E+01 kg Renewable material resources 8.3E+00 Renewable primary energy MJ 4.5E-01 Consumption of freshwater $m^3$ 7.7E-04

3. Material composition				
Material		Unit		
Shou sugi ban	100	%		
Package (film)	0.021	%		
Hotmelt	0.090	%		

4. Waste to disposal		
Parameter		Unit
Hazardous waste	0.0E+00	kg
Non-hazardous waste.	4.5E-04	kg

<sup>\*</sup>Data derived from LCA and not assigned to the impact categories of LCIA



For the analysis, a set of foreground data was first prepared based on the foreground data collected for one year (October 2017 to September 2018) and then they were multiplied by the pertinent background data to estimate environmental loads. Transportation was calculated by collecting actual data over one year. As the product is manufactured in the plants in Hiroshima and Tokushima Prefectures, the averages of data taken from the two plants were used to represent the product data.

The analysis revealed that dominant stages varied depending on the LCI parameters (see the graph above). Namely, Ozone layer destruction, Eutrophication were under the stronger influence of A1 Raw Material Supply, while Global warming, Photochemical ozone were predominantly affected by A3 Manufacturing.

The carbon storage was calculated based on Annex F of the PCR as follows:

Carbon Storage (kg-C) = 6.06 (kg-wood)  $\times 0.5$  = 3.03 (kg-C) (=11.1kg-CO2)

### 6-1. Supplementary environmental information



**Ecoleaf Environmental Labeling Program** 

Sustainable Management Promotion Organization Type III Environmental Declaration (EPD) 2-1, Kaji-cho 2 chome, Chiyoda-ku, Tokyo Japan Registration number: JR-BC-20002E

https://ecoleaf-label.jp/

6-2. Regulated hazardous substances					
Substance	CAS No.	Reference to standards or regulations			
-					

### 7. Assumptions of secondary data used

Inventory Database: IDEA Ver.2.1.3

### 8. Remarks

- For data quantification, please refer to PCR and Rules on quantification and declaration.
- Comparative assertion is permitted only when Rules on quantification and declaration are satisfied. (Reference URL: https://ecoleaf-label.jp/regulation/)

Registration number: JR-BC-20002E